
# Top Quark Production Asymmetry at the Tevatron

Dan Amidei University of Michigan

# top A<sub>FR</sub> in NLO QCD



- $C = -1 + C = +1 \rightarrow A_C$
- at Tevatron this is an A<sub>FB</sub>

• measure in 
$$\Delta y = y_t - y_{\bar{t}}$$

$$A^{t\bar{t}} = \frac{N(\Delta y > 0) - N(\Delta y < 0)}{N(\Delta y > 0) + N(\Delta y < 0)}$$

inclusive NLO prediction (QCD+EWK)

$$A_{FB} = (6.6 \pm 2.0 + 2.0?)\%$$

measured

$$A_{FB} \sim (19 \pm 4)\%$$

- new work on SM calculation
  - EWK, denominators
  - t-tbar P<sub>t</sub> parton showers
  - NNLO is coming
- many interesting NP models
  - s-channel, t-channel

# top ∆y asymmetry in I+jets

#### similar analyses


- subtract backgrounds
- kinematic reco of top 4-vectors
- SVD unfold


#### CDF

- arXiv:1211.1003
- 9.4 fb<sup>-1</sup>, 2653 evnts 530 bkg
- differential xsec in  $\Delta y$
- $-A_{fb}$  parton =  $(16.2 \pm 4.7)\%$

#### • D0


- PRD84, 112005 (2011)
- 5.4 fb-1, 1581 evnts 455 bkg
- $A_{fb}$  parton =  $(19.6 \pm 6.5)\%$





# comparison of tevatron top $\Delta y A_{FB}$ results Sept. 2012


 including private average simple weighted, neglect correlations




• expected precision w 9 fb<sup>-1</sup> for all measurements < 3%.

- rapidity dependence
  - $\text{ fit } \chi^2_{\text{p.d.f.}} = 1.0$
  - slope >  $3\sigma$  from 0
  - PE how often  $\alpha_{NLO}$  ≥  $\alpha_{data}$
  - use background sub data
  - $p_{NLO} = 0.00892$

- mass dependence
  - slope is  $>3\sigma$  from 0
  - fit  $\chi^2_{p.d.f.} = 0.3$
  - $p_{NLO} = 0.00646$






### differential top cross-section in production angle $\cos \theta^*$


#### full reconstruction in tt frame

- study production angle directly
- SM  $\sim 1 + \cos^2\theta$
- NLO adds asymmetry
- s-channel models add cosθ term
- t-channel leading term  $1/(1-\cos\theta)$
- some benchmark models on right

#### new CDF analysis in I+jets

- 9.4 fb<sup>-1</sup> 3864 evnts, 1026 bkg
- reconstruct  $cos(\theta^*)$
- decompose in 8 Legendre moments
- no binning, no regularization
- estimate moments response from MC
- invert response, multiply by bkg subtracted moments
- integrate series over 10 bins






### study the scattering angle directly

#### moments

- SM agreement except L = 1
- $a_1 = 0.44 \pm 0.12$  measured 0.15±.05 expected
- excess linear term in xsec
- benchmark Z' model is disfavored

- contribution of moments to A<sub>fb</sub>
  - independent asymmtries add!
  - A<sub>FB</sub> is entirely due to linear term



# single lepton in I+jets

#### lepton

- well measured
- follows top quark
- A(qy<sub>I</sub>) robust asymmetry probe
- correlation with A(∆y) sensitive to polarization

#### pioneered by D0


- PRD84, 112005 (2011)
- 5.4 fb-1, 1581 evnts 455 bkg


#### data A<sub>fb</sub>

- $(14.2 \pm 3.8)\%$  obs
- (0.8 ± 0.6)% pred

#### parton A<sub>fb</sub>

- $(15.2 \pm 4.0)\%$  obs
- (2.1 ± 0.1)% pred





# D0 dilepton rapidity asymmetries


- arXiv:1207.0364
- 5.4 fb<sup>-1</sup>, 2 OS leptons (M.ne.Z) + met + 2 jets + Ht
- $649 \text{ events bkg} = 244 \pm 18$
- no tt reconstruction: it's the leptons


				1
	Raw	Unfolded	Predicted	•
$A^{\ell}$	$2.9 \pm 6.1 \pm 0.9$	$2.5 \pm 7.1 \pm 1.4$	$4.7 \pm 0.1$	$A_{\mathbb{C}}$
$A_{ m FB}^{\ell^+}$	$4.5 \pm 6.1 \pm 1.1$	$4.1 \pm 6.8 \pm 1.1$	$4.4 \pm 0.2$	+ charge leptons
$A_{\mathrm{FB}}^{\ell^-}$	$-1.2 \pm 6.1 \pm 1.3$	$-8.4 \pm 7.4 \pm 2.4$	$-5.0 \pm 0.2$	- charge leptons
$A_{\mathrm{FB}}^{\ell}$	$3.1 \pm 4.3 \pm 0.8$	$5.8 \pm 5.1 \pm 1.3$	$4.7 \pm 0.1$	both charges (q.η)
$A^{\ell\ell}$	$3.3 \pm 6.0 \pm 1.1$	$5.3 \pm 7.9 \pm 2.9$	$6.2 \pm 0.2$	Δy leptons
$A_{\mathrm{CP}}^{\ell}$	$1.8 \pm 4.3 \pm 1.0$	$-1.8 \pm 5.1 \pm 1.6$	$-0.3 \pm 0.1$	CP violating

# inclusive lepton asymmetry at CDF

- study q·y<sub>I</sub>
  - "asymmetric part" A<sub>I</sub>(qy<sub>I</sub>)
  - symmetric part is model indpndnt
- ansatz  $\mathcal{A}(qy_l) = a \tanh \left[\frac{1}{2}qy_l\right]$ 
  - well fit in benchmark models
    - OctetA heavy, unpolarized
    - OctetsLR light, polarized
  - extrapolates into forward region
- technique
  - integrate with symmetric part
  - gets full distribution + total asymmetry
  - works well in benchmarks
- measurement
  - 9.4 fb<sup>-1</sup> 3864 evnts, 1026 bkg

$$A = 0.094 \pm 0.024^{+0.022}_{-0.018}$$





#### combined lepton asymmetry

D0 combination

$$-A_{FB}^{I}DIL = (5.8 \pm 5.3) \%$$
 pred  $(4.7 \pm 0.1)\%$ 

- $-A_{FB}^{I}$  I+jets = (15.2 ± 4.0)% pred (2.1 ± 0.1)%
- D0 combination  $A_{FB}^{I} = (11.8 \pm 3.2)\%$
- CDF I+jets

$$-A_{FB}^{I}$$
 I+jets =  $(9.4 \pm 3.1)$  % pred  $(3.8 \pm 0.3)$ %


- informal combination
  - $-A_{FB}^{I}$  = (10.5 ± 2.2)% exceeds SM predictions by 3-4  $\sigma$
- what do we expect for SM decays with measured A(Δy)?
- crude measure

$$\frac{A_{FB}^{l}}{A_{FB}^{\Delta y}}\Big|_{powheg} \times A_{FB}^{\Delta y}\Big|_{data} = 0.46 \times (18.7 \pm 3.7) = 8.6 \pm 1.7$$

would be interesting to have a real prediction


# p<sub>t</sub> (tt) dependence of the asymmetry at CDF

- examine at obs bkg-sub level
  - pythia and powheg follow expected trend
  - CDF 8.7 fb<sup>-1</sup> l+jets
  - data above predictions
- normalize predictions to the data
  - independent asymmetries add!
  - scale each bin by inclusive A<sub>fb</sub>
- good agreement with either Powheg and Pythia
- excess asymmetry is P<sub>t</sub> independent?



# p<sub>t</sub> (tt) dependence of the asymmetry at CDF

- examine at obs bkg-sub level
  - pythia and powheg follow expected trend
  - CDF 8.7 fb<sup>-1</sup> l+jets
  - data above predictions
- normalize predictions to the data
  - independent asymmetries add!
  - scale each bin by inclusive A<sub>fb</sub>
- good agreement with either Powheg and Pythia
- excess asymmetry is P<sub>t</sub> independent?



#### summary

- inclusive asymmetry in agreement CDF+D0
  - informal combo  $A^{tt}_{FB} \sim (18.7 \pm 3.7)\%$
  - eventual combined  $\delta A_{FB} \sim 3.0\%$
- linear  $M_{tt}$  and  $\Delta y$  dependence of  $A_{fb}$  in tt system (CDF)
  - slopes  $3\sigma$  from zero and  $2\sigma$  larger than NLO prediction
- dσ/dcos(θ\*) (CDF)
  - decompose in legendre moments
  - excess linear term
- measured 2-3 $\sigma$  asymmetry in the lepton alone (D0 +CDF)
  - slightly high for measured  $A(\Delta y)$ ?
- p<sub>t</sub>(tt) dependence agrees with Poweg/Pythia + p<sub>t</sub> independent offset
- some tension between D0 dileptons and everything else
- picture still incomplete, much work still to do

additional material

# lepton asymmetry method

$$\mathcal{S}(qy_{l}) = \frac{N(qy_{l}) + N(-qy_{l})}{2}$$

$$\mathcal{A}(qy_{l}) = \frac{N(qy_{l}) - N(-qy_{l})}{N(qy_{l}) + N(-qy_{l})}$$


$$N(qy_{l} > 0) = \int_{0}^{\infty} dqy_{l} \left[ \mathcal{S}(qy_{l}) \times (1 + \mathcal{A}(qy_{l})) \right]$$

$$N(qy_{l} < 0) = \int_{0}^{\infty} dqy_{l} \left[ \mathcal{S}(qy_{l}) \times (1 - \mathcal{A}(qy_{l})) \right]$$

$$A_{FB}^{lep} = \frac{N(qy_{l} > 0) - N(qy_{l} < 0)}{N(qy_{l} > 0) + N(qy_{l} < 0)}$$

$$= \frac{\int_{0}^{\infty} dqy_{l} \left[ \mathcal{A}(qy_{l}) \times \mathcal{S}(qy_{l}) \right]}{\int_{0}^{\infty} dqy_{l} \mathcal{S}(qy_{l})}$$

# lepton asymmetry method



### lepton asymmetries

- lepton follows top
  - independent of asymmetry mechanism if P=0
- reconstructed lepton η is systematically unencumbered
- "bias free" asymmetry indicator

#### single lepton

# $A_{FB}^{l^{\pm}} = \frac{N_{l^{\pm}}(\eta > 0) - N_{l^{\pm}}(\eta < 0)}{N_{l^{\pm}}(\eta > 0) + N_{l^{\pm}}(\eta < 0)}$


$$A_{FB}^{l} = \frac{N_{l}(Q \cdot \eta > 0) - N_{l}(Q \cdot \eta < 0)}{N_{l}(Q \cdot \eta > 0) + N_{l}(Q \cdot \eta < 0)} \qquad A_{CP}^{l} = \frac{N_{l^{+}}(\Delta \eta > 0) - N_{l^{-}}(\Delta \eta < 0)}{N_{l^{+}}(\Delta \eta > 0) + N_{l^{-}}(\Delta \eta < 0)}$$

#### two lepton


$$A^{ll} = \frac{N(\Delta \eta > 0) - N(\Delta \eta < 0)}{N(\Delta \eta > 0) + N(\Delta \eta < 0)}$$

$$A_{CP}^{l} = \frac{N_{l^{+}}(\Delta \eta > 0) - N_{l^{-}}(\Delta \eta < 0)}{N_{l^{+}}(\Delta \eta > 0) + N_{l^{-}}(\Delta \eta < 0)}$$

# lepton A<sub>FB</sub> performance in the W+1 jet sample (CDF)



	$ \eta_{lep}  < 0.75$	$ \eta_{lep}  \ge 0.75$
Observed Data	$0.059 \pm 0.001$	$0.124 \pm 0.002$
SM Prediction	$0.063 \pm 0.005$	$0.134 \pm 0.008$
Data Minus Prediction	$-0.004 \pm 0.005$	$-0.010 \pm 0.008$



	$p_T < 60 \text{GeV}/c$	$p_T \ge 60 { m GeV}/c$
Observed Data	$0.083 \pm 0.001$	$-0.009 \pm 0.004$
SM Prediction	$0.089 \pm 0.004$	$-0.001 \pm 0.013$
Data Minus Prediction	$-0.006 \pm 0.004$	$-0.008 \pm 0.014$

# Historical perspective

